

Exercises for Further Graphics (Lectures 1-4)

All work to be submitted by email in a single PDF, no less than 48 hours before supervision.

1. Terms and Concepts

Voronoi Diagrams

- What is *equiangularity*?
- What is the *empty circle property*?
- Describe how to use hardware acceleration to swiftly compute Voronoi diagrams. What are the limitations of this approach?

Topology

- Define the *Euler characteristic*
- Define the term *angle deficit*
- State the *Poincaré Theorem*, which links the geometry of a surface to its topology
- State Descartes' *Theorem of Total Angle Deficit*, which links angle deficit across a surface to its Euler characteristic

Curvature

- The *one-ring* of a vertex is the (usually ordered) set of vertices which lie exactly one edge away from a given vertex on a polyhedral surface. Given a vertex V with one-ring $\{v_0, \dots, v_{n-1}\}$, give a formula for the discrete curvature of the surface at V .

2. Signed Distance Functions

Give signed distance functions for:

- A cone
- An igloo
- An arbitrary tetrahedron
- The spiral of a corkscrew

3. Bezier curves

- Why is a Bezier curve contained entirely within the convex hull of its control points?
- Give real-world examples of C0, C1, C2 continuity
- Prove that the linear interpolation of two linear interpolations is, in fact, a Bezier quadratic.

4. B-Splines

- a. Show that the B-spline with $k = 3$ and knot vector $[0 \ 0 \ 0 \ 1 \ 1 \ 1]$ is equivalent to the quadratic Bezier curve.
- b. Give a knot vector and value of k which would describe a uniform B-spline equivalent to a cubic Bezier curve.
- c. Derive the formula of and sketch a graph of $N_{3,3}(t)$, the third of the quadratic B-spline basis functions, for the knot vector $[0 \ 0 \ 0 \ 1 \ 3 \ 3 \ 4 \ 5 \ 5 \ 5]$.

5. Barycentric coordinates

What does it mean if one or more of the coefficients of the barycentric coordinates of a point with respect to a triangle are negative?